

Processing Solid Waste of Animal Beef using Bioactivators in Effort to Reduce Environmental Pollution

by Titik Mariyati Indasah

Submission date: 14-Apr-2022 03:42PM (UTC+0700)

Submission ID: 1810479080

File name: PH_VOL_5_NOV_21_TITIK_MARYATI_TERAKREDITASI_SINTA_5_NILAI_15.pdf (379.58K)

Word count: 5194

Character count: 26539

Processing Solid Waste of Animal Beef using Bioactivators in Effort to Reduce Environmental Pollution

Titik Mariyati¹, Indasah²

¹ SMK Kesehatan Bhakti Indonesia Medika Kediri
² Institute of Health Sciences Strada Indonesia

Email:

titik.mariyati@ymail.com

ABSTRACT

Cattle waste both in the form of solid and liquid waste that is produced every day if left alone without proper waste treatment, it can cause environmental pollution, namely air, water, and soil pollution. The environment becomes squalid and attracts various animals such as flies and dogs which can carry various diseases such as deare, typhoid, cholera and skin diseases which can spread quickly and smell bad. One of the efforts to overcome environmental pollution around cattle is to carry out a composting process using cow rumen bioactivator. This study aims: (1) to find out whether cow rumen bioactivator can reduce the unpleasant odor caused by cattle waste, (2) find out whether cow rumen bioactivator can accelerate the composting process, (3) determine the effect of bioactivator volume on compost quality, namely nitrogen content. (N-total), Phosphorus levels, and Potassium levels. This study uses qualitative analysis to test odors (H₂S gas, CO₂ gas, NH₃ gas and CH₄ gas) which was carried out at the CHEMICAL Laboratory of SMK Kesehatan Bhakti Indonesia Medika Kediri, and quantitative analysis to test Nitrogen levels using a Kjeldahl flask, Phosphorus levels with a visible spectrophotometer. , Potassium levels with AAS conducted at the Laboratory of PTPN X Jengkol Plosoklaten Kediri. This study showed: (1) after the composting process, cattle solid waste did not smell, (2) the composting process took 30 days, (3) the addition of various bioactivators (25 ml, 50 ml, 75 ml, and 100 ml), volume 50 ml is the optimal volume in the composting process with 3 kg of cattle solid waste. From the decomposition results, total nitrogen content was 0.592%, phosphorus content was 0.27%, and potassium content was 0.538%. These levels meet the Quality Standards for organic fertilizers and soil enhancers according to the Regulation of the Minister of Agriculture no. 2/Pert./HK.060/2/2006.

Received : October 4nd 2021

Accepted : October 15rd 2021

Published : November 27th 2021

Keywords: Environmental pollution, cattle solid waste, cow rumen bioactivator, variations of bioactivator

Copyright © 2021 IIK STRADA Indonesia
All right reserved.



¹ This is an open-access article distributed under the terms of the Creative Commons Attribution-ShareAlike 4.0 International License.

INTRODUCTION

Livestock waste can be in the form of solid (faeces), liquid (urine), and gas (H₂S, NH₃, CO₂ and CH₄). Untreated livestock waste will be a source of pollutants for water, air, and soil pollution.

As many as 55% of the people in Jimbir Sugihwaras Prambon Nganjuk village are cattle and goats. The livestock waste produced consists of solid waste in the form of animal feces and feed residues, as well as liquid waste in the form of cage washing wastewater, livestock sanitation wastewater and livestock urine. If not handled properly, the livestock waste has the potential to cause environmental pollution and health problems for the community around the farm.

Cattle that have to produce solid, liquid and gas waste every day. The solid waste of cattle produced in rearing activities has been left untreated, left alone, it will make the solid waste of cattle accumulate more and more and there is no special place used to accommodate the waste, it will be a complicated problem for the environment around the barn. This is what causes a dirty environment and becomes a breeding ground for harmful microorganisms such as worms or flies. In addition, the solid waste of cattle also causes an unpleasant / pungent odor around the cattle pens. Generally, livestock manure is thrown away or simply placed around the cage environment without any effective treatment, this will cause the cage environment to become dirty, and in months if it is not immediately handled to become compost, the livestock manure will cause environmental pollution in the form of air pollution from the unpleasant odor caused by the manure, and can also cause the environment to become slum, and attractive to various animals, such as flies and dogs, which can carry disease. Potential health hazards that can be caused such as diarrhea, cholera, typhoid and mold can spread quickly, because improperly managed waste can mix with drinking water and spread disease viruses. (Hidayat, 2017). Thus, to keep the environment around cattle farmers clean and healthy, it is necessary to process cattle solid waste into solid organic fertilizer in a relatively short time of 3 to 4 weeks with the addition of a bioactivator.

Bioactivator, also known as Effective Microorganism (EM) is a collection of various fermentative microbes, which function in the fermentation of organic materials. There are five main groups of fermentative microbes including: photosynthetic bacteria, Lactobacillus, Streptomyces, yeast (yeast), and Actinomycetes (Wiryanti, 2014). From this problem, it is hoped that the processing of cattle solid waste with the addition of a bioactivator is expected to reduce environmental pollution

Processing of cattle manure needs to be done to reduce environmental pollution, by treating cattle waste as manure because it contains nutrients such as nitrogen (N), phosphorus (P), and potassium (K) as well as micro nutrients including calcium, magnesium, sulfur, sodium, iron, and copper needed by plants and soil fertility (Hapsari, 2013) Therefore, before being disposed of or used as fertilizer or soil enhancer, these materials must be stabilized first so as not to negatively affect the soil and plants.

MATERIALS AND METHODS

Types of research

The type of research used in this research is real research (True Experiment). Using True Experiment (real research) because the subject group (test substance) was chosen randomly, there was a comparison group to the group being treated, and there was control over the conditions in order to minimize the influence of other variables (interfering).

According to Sumadi Suryabrata (1990: 32-36) True Experiment aims to investigate the possibility of a causal relationship with the design where in fact there is a treatment group and a control group that is not subject to treatment conditions.

Research design

Research design is the methods used to collect research data so that research can be proven. In this study, the "pretest-posttest control groups design" was used, namely there were two groups selected at random, then given a pretest to determine whether there was a difference between the experimental group and the control group in the initial state. Pretest results are good if the experimental group scores are not significantly different.

The research design used in this study was a "pretest-posttest control groups design", namely a research design that contained a pretest before being given treatment and a posttest after being given treatment. Thus it can be known more accurately, because it can be compared with being held before being given treatment. (Sugiyono, 2001:64),

12

Table of Research Design Pretest Posttest Control Group Design


Kelompok	Pretest	Perlakuan	Posttest
E	P1	X	P2
K	P3	-	P4

Information :

- E : Experimental Group (the group that was treated with the addition of bioactivator)
- K : Control group (the group that was not treated with the addition of bioactivator)
- P1 : Pretest experimental group
- P2 : Posttest experimental group
- P3 : Pretest control group
- P4 : Posttest control group
- X : Treatment with addition of bioactivator

27

Image of the pretest-posttest control groups design:



Information :

- Sample : Test Substance (bovine solid waste)
- Pretest : Initial test
- MSOP : Matched Subject Ordinal Pairing
- E : Experimental Group
- K : control group
- X : Treatment with addition of bioactivator
- T : Without the addition of bioactivator treatment
- Posttest : Final test

The composting result from this experiment were compared with the Quality Standards for organic fertilizers and soil enhancers according to the Regulation of the Minister of Agriculture No. 2/Pert./HK.060/2/2006

Table 1. Minimum Technical Requirements for Organic Fertilizer

No	Requirements	Unit	Parameters	
			solid	liquid
1	C-Organic	%	> 12	≥ 4,5
2	C/N ratio	%	10 - 25	
3	Follow-up materials (gravel, glass, plastic, etc.)	%	Maks 2	
4	Granule	%	4 - 12	
	bulk		13 - 20	
5	Heavy metal content			
	As	ppm	≤ 10	≤ 10
	Hg	ppm	≤ 1	≤ 1
	Pb	ppm	≤ 50	≤ 50
	Cd	ppm	≤ 10	≤ 10
6	PH		4 - 8	4 - 8
7	Total rate			
	P2O5	%	< 5	< 5
	K2O	%	< 5	< 5
8	Micro Elements	%		5
	Zn		Max 0,5	Max 0,25
	Cu		Max 0,5	Max 0,25
	Mn		Max 0,5	Max 0,25
	Co		Max 0,002	Max 0,0005
	B		Max 0,25	Max 0,125
	Mo		Max 0,001	Max 0,001
	Fe		Max 0,4	Max 0,004

Sumber : Direktorat Sarana Produksi, 2006

1

Table 2. Nutrient and Water Content of Several Types of Manure

Type of Livestock	Nutrient and Water Content (%)				Information
	Nitrogen	Information	Potassium	Water	
Cow					
Congested	0,40	0,20	0,10	85	Cold fertilizer
Liquid	1,00	0,50	1,50	92	
Buffalo					
Congested	0,60	0,30	0,34	85	Cold fertilizer
Liquid	1,00	0,15	1,50	92	
Goat					
Congested	0,60	0,30	0,17	60	Hot fertilizer
Liquid	1,50	0,13	1,80	85	
Sheep					
Congested	0,75	0,50	0,45	60	Hot fertilizer
Liquid	1,35	0,05	2,10	85	
Chicke	1,00	0,80	0,40	55	Cold fertilizer

Sumber : Ade Iwan Setiawan, 2007

Research Location and Time

Locations The research was conducted at:

- The fermentation process is carried out in a cowshed in Jimbir Hamlet, RT 02 RW 08 Sugiharwas, Prambon District, Nganjuk Regency.
- Tests for Nitrogen, Phosphorus, and Potassium levels were carried out at the Sugar Research Center Laboratory of PTPN X in Plosoklaten Kediri.
- Testing the qualitative analysis at the CHEMICAL Laboratory of SMK HEALTH BHAKTI INDONESIA MEDIKA KEDIRI

Research time

The research was carried out for 2 months, starting from May 1, 2021 to June 30, 2021.

Tools and materials

The tools and materials in the research consisted of 2 kinds, namely:

- Tools and materials used in the cage as an experiment (composter)
 - Equipment in the fermentation process
Tong, gloves, measuring cup, mask, plastic bag, bucket, hoe, arco, water hose, scale (balance), thermometer, scissor, spray bottle, raffia, elastic rubber, book, pen.
 - Ingredients in the fermentation process
Fresh cow dung (rumen), molasses, coconut water, rice bran, rotting fruit residue, solid cow dung, well water.
- Tools and materials for the analysis of compost quality parameters. at the Research Center Laboratory of PTPN X Plosoklaten Kediri.
 - Tools for analyzing compost quality parameters
Analytical balance with 0.1 mg accuracy, degeston apparatus (electric heater), distillation unit, Kjeldahl flask, burette, dispenser, Erlenmeyer flask, volume pipette, 50 ml or 100 ml volumetric flask, Digestion flask, microwave digester, 20 volume chemical tube ml, Vortex mixer, Visible Spectrophotometer, AAS (Atomic Absorbance Spectrophotometer), Cathode lamp K.
 - Materials on compost quality parameter analysis
Water, concentrated H₂SO₄, sulfuric-salicylic acid solution (dissolved 25 g of salicylic acid in 1 liter of concentrated sulfuric acid, sodium thiosulfate, standar solution of H₂SO₄ 0.05 N, Boric acid 1% w/v, Conway indicator (added 0.15 g bromo cresol green + 0.1 g methyl red in 100 ml alcohol 96 %), NaOH 40 %, HNO₃ 65%, HClO₄ 70 %, KH₂PO₄, Phosphate Molybdate reagent, LaCl₃

Research Implementation Design

This study used a randomized design consisting of 20 formulations of compost material to be treated, as follows:

- P0 = 3 kg of solid cow dung (normal)

P1 = 3 kg cow dung + 25 ml cow rumen bioactivator

P2 = 3 kg cow dung + 50 ml cow rumen bioactivator

P3 = 3 kg cow dung + 75 ml cow rumen bioactivator

P4 = 3 kg cow dung + 100 ml cow rumen bioactivator

Each treatment was repeated 5 times, so that 20 experimental plastic bags were obtained

Research Stages

The research was conducted through 3 stages, namely:

1. Introduction Stage

The introductory stage is the stage that begins with the collection of cow rumen and cow solid manure

2. Experimental stage

The experimental stage is divided into 2, namely:

2.1. Stages of making bioactivator

a. 10 kg of beef rumen plus 1 liter of molasses later

b. plus 5 kg of rice bran which is then diluted with enough coconut water, then

c. placed in a barrel, allowed to ferment for 1 week.

d. After fermentation occurs, it is filtered, then the filtrate is added to the rest of the rotten fruit.

e. The bioactivator is ready to use.

2.2. The implementation stage of the research was to put 3 kg of solid cow dung into a plastic bag then add a variety of bioactivators from the cow's rumen and put it in a plastic bag and then put the bucket in a closed room.

3. Decomposition analysis stage

The decomposition analysis stage is that during the decomposition process until the composting process is complete, several measurements are carried out once a week including measurements of temperature, pH, color and odor. After the compost is formed, it is continued by analyzing the levels of Nitrogen, Phosphorus, and Potassium, and qualitative analysis of odor.

Sampling Technique

Sampling (test substance) using purposive sampling technique, which means the test substance is taken from cow dung that one day comes out of the stomach of cattle.

Research variable

1. Independent variable

The independent variables in this study include: Bioactivator volume

2. Bound Variable

The dependent variables in this study include: the weight of the test substance (bovine solid waste), and the quality of the compost

Data analysis

1. Data Analysis Requirements

a. Data Distribution Normality Test

Normality test aims to determine whether a research data is normally distributed or not. In parametric statistical analysis, data with normal distribution is a must as well as an absolute requirement that must be fulfilled. If the data is not normally distributed, then the research hypothesis is tested using a non-parametric statistical analysis approach.

One way to detect the normality of a data can be done with the Kruskal-Wallis technique. The Kruskal-Wallis test is called the H₁₅ test, which is an alternative procedure to the one-way ANOVA. Kruskal-Wallis is a non-parametric alternative method that can be used for ordinal response data or ranked data.

b. Data Homogeneity Test

The homogeneity test aims to determine whether the variation of some data from the population has the same variance or not. This homogeneity test serves as a

requirement (though not an absolute requirement) in comparative analysis such as the independent sample t test and ANOVA test.

According to Joko Widiyanto (2010:51) the basis or guidelines for decision making the homogeneity test are as follows:

1. If the significance value or Sig. <0.05, it is said that the variance of two or more population groups is not the same (not homogeneous).
2. If the significance value or Sig. > 0.05, it is said that the variance of two or more data population groups is the same (homogeneous).

2. Hypothesis test

The data analysis technique used in this research is ANOVA analysis. This technique aims to determine the difference between the actualization of the composting process in the application of a bioactivator and the composting process without the addition of a bioactivator. The hypotheses that the researcher proposes are:

- a. H1 or the first hypothesis: There is an effect of adding bioactivator to the time used in the composting process of cattle solid waste.
- b. H2 or second hypothesis: There is an effect of adding a bioactivator to the smell of cattle solid waste.
- c. H3 or third hypothesis: There is an effect of adding bioactivator to compost quality (content of N, P, K elements in cattle solid waste).

To test the research hypothesis above, we need to make a decision in the ANOVA test, the decision that can be obtained is by looking at the significance value (sig), based on the significance value (Sig)

1. If the significance value (Sig) < probability 0.05, then there is an effect of the independent variable on the dependent variable, or the hypothesis is accepted.
2. If the significance value (Sig) > 0.05 probability, then there is no effect of the independent variable on the dependent variable, or the hypothesis is rejected.

RESULTS

In this section, the researcher will convey the results of the research which include:

- A. Qualitative analysis stage
- B. Decomposition analysis stage

A. Qualitative Analysis Stage

Examination Results Qualitative analysis.

No	Zat Uji	H2S	CO2	NH3	CH4	PH	Smell	color
1	PB	+	+	+	+	6,8	Not delicious and pungent	yellowish
2	PL	+	+	+	-	6,8	Not delicious	chocolate
3	P1	-	-	-	-	6,8	-	Dark chocolate
4	P2	-	-	-	-	6,8	-	Dark chocolate
5	P3	-	-	-	-	6,8	-	Dark chocolate
6	P4	-	-	-	-	6,8	-	Dark chocolate

B. Decomposition Analysis Stage

Compost Quality Measurement Results (N, P, K)

Replication	Treatment	Content	Bioactivator Volume (%)					
			PB	PL	P1	P2	P3	P4
I	N		0.31	0.58	0.58	0.58	0.55	0.59
	P		0.18	0.33	0.10	0.60	0.14	0.33
	K		0.31	0.44	0.57	0.58	0.33	0.44
II	N		0.31	0.58	0.56	0.6	0.61	0.45
	P		0.18	0.33	0.32	0.15	0.20	0.25

	K	0.31	0.44	0.52	0.52	0.37	0.38
III	N	0.31	0.58	0.58	0.60	0.60	0.59
	P	0.18	0.33	0.25	0.20	0.22	0.30
	K	0.31	0.44	0.55	0.56	0.56	0.58
IV	N	0.31	0.58	0.58	0.58	0.59	0.59
	P	0.18	0.33	0.20	0.15	0.21	0.22
	K	0.31	0.44	0.50	0.56	0.57	0.49
V	N	0.31	0.58	0.32	0.60	0.61	0.60
	P	0.18	0.33	0.32	0.25	0.30	0.30
	K	0.31	0.44	0.54	0.57	0.58	0.58

Information :

PB: Test substance that just came out of cattle

PL : Test substance without treatment left for 4 weeks

P1 : The test substance is treated with the addition of 25 ml of bioactivator

P2 : The test substance is treated with the addition of 50 ml of bioactivator

P3 : Test substance with the addition of 75 ml of bioactivator

P4 : The test substance is treated with the addition of 100 ml of bioactivator

DISCUSSION

In the compost maturation process, the degree of acidity ranges from PH 6 to 7 (neutral PH). The pH of the compost serves as an indicator of the compost decomposition process. Compost microorganisms will work to carry out their activities and grow at a PH between 5.5 to 8. (PINLITAMAS 1: 2018

In the composting process, temperature is a very important factor in determining the proliferation of microorganisms. Microorganisms are bacteria that will decompose solid waste cow manure to become compost. At the beginning of composting the temperature of the test substance was 27oC.

After an interval of 1 week (day 8) the temperature began to rise, the test substance without treatment (control) rose to 28°C, while those with treatment increased significantly by 33 °C and 35 °C. In this phase, microorganisms are present and begin to work on decomposing cattle solid waste.

In the second week (16th day) for the test substance without treatment (control) the temperature remained at 28 °C, while the test substance with temperature treatment increased dramatically, here is the peak temperature of composting, i.e. some are at a temperature of 35°C, some are a temperature of 36 °C and there is a temperature of 37 °C.

In the third week (day 21) for the test substance without treatment (control) the temperature remained at 28 °C, while for the test substance that was being treated the temperature began to drop, namely at a temperature of 30°C, 32 °C and 32 °C. In this phase the activity of microbes (microorganisms) begins to decrease. This phase is known as the compost maturation phase.


On the fourth week (day 30) all the test substances were at 28 °C which is at room temperature. (Article : Chem.Notes 2020)

The temperature in the composting process gradually decreases due to reduced organic matter that can be decomposed by microorganisms. This can occur because the life of mesophilic and thermophilic microorganisms alternately. When t₂₂ temperature drops, mesophilic microorganisms grow to replace thermophilic microorganisms. (Journal.unnes.ac.id/sju/index.php/Unnes.JLifeSci, 2016)

In the anaerobic composting process when added microorganisms (Bioactivator) it will last 30 days. (*Artikel : Perencanaan, Perancangan Lingkungan 2021*)

Nitrogen Content

Figure From the decomposition process, the data obtained on the average levels of N, P, K are as :




In the Nitrogen content histogram above, the standard Nitrogen content is 0.4%. Nitrogen content in cattle solid waste that just came out of cattle 0.31% does not meet the standard of use.

Meanwhile, cattle solid waste that was left for 30 days without treatment had a concentration of 0.58% and that which was treated with added bioactivator met the appropriate standard for use because the level was already above 0.4%. (*Setiawan, Ade Iwan. 2007*)

Cattle solid waste that is left for 30 days and added with a bioactivator, the nitrogen content increases, this is because there is already a role for microorganisms that decompose cow solid waste.

Nitrogen levels from cattle solid waste that are left for 30 days without treatment are higher than nitrogen levels from livestock solid waste added with bioactivators, because in the decomposition process microorganisms also need nitrogen elements to maintain and form body cells. In addition, nitrogen can react with water to form NO_3^- and H^+ . Where NO_3^- can occur decomposition reactions into N_2 gas and N_2O gas. (*Journal of Veterinary Science (1), June 2017*)

Phosphorus Level




In the Phosphorus histogram above, the standard Phosphorus level is 0.2% suitable for use. Phosphorus levels in cattle solid waste that just came out of cattle 0.18% do not meet the standard. Meanwhile, cattle waste that was left for 30 days without treatment had a concentration of 0.33% and that which was treated with added bioactivator met the proper use standard because the level was already above 0.2%. (*Setiawan, Ade Iwan. 2007*), (*No.2/Pert.IHK.060/2/2006*)

By leaving it for 30 days and adding a bioactivator, the Phosphorus content increases, this is because there is already a role for microorganisms that decompose solid cattle waste.

Phosphorus levels from cattle solid waste that were left for 30 days without treatment were higher than the levels of Phosphorus from livestock solid waste added with bioactivators, because in the process of decomposing cattle solid waste microorganisms in their activities also require Phosphorus elements. (*ISSN : 2621 – 1475, J.Fish Protech 2020, vol.3 No.2*)

Potassium Level



In the histogram of Potassium levels above, the standard Potassium level is 0.1% suitable for use. Potassium levels in cattle solid waste that just came out of cattle 0.31% meet the standard of use. (Setiawan, Ade Iwan. 2007), (No.2/Pert./HK.060/2/2006)

Meanwhile, cattle solid waste that was left for 30 days without treatment had a concentration of 0.44% and that which was treated with added bioactivator met the proper use standards because the levels were already above 0.1%.

Cattle solid waste that is left for 30 days and added with a bioactivator, the content of Potassium increases, this is because there is a role for microorganisms that decompose solid cattle waste.

Potassium levels from cattle solid waste that were left for 30 days without treatment were lower than Potassium levels from livestock solid waste added with bioactivator. Thus, the addition of a bioactivator greatly affects the composting process. Elemental Potassium is used by microorganisms in the substrate material as a catalyst, so that an increase in the activity of microorganisms will increase the level of Potassium. (Journal of Veterinary Science (1), June 2017)

8

CONCLUSION

Based on the results of research and discussion, the following conclusions can be drawn:

1. The addition of a bioactivator can accelerate the composting process of cattle solid waste for 4 weeks (30 days), so as to create a clean and healthy environment.
2. The addition of a bioactivator can reduce unpleasant and pungent odors in cattle solid waste because NH₃, H₂S, CO₂ and CH₄ gases have been decomposed into other elements (N, C) within 30 days, so as to reduce air pollution.
3. The optimal addition of a bioactivator in the processing of cattle solid waste based on the quality of the compost is the addition of a 50 ml bioactivator.
4. The quality of the compost (concentration of N, P, K) has met the standards of the Regulation of the Minister of Agriculture No.2/Pert./HK.060/2/2006.
5. Based on statistical tests, then:
 - a. The addition of bioactivator affects the formation of nitrogen nutrients.
 - b. The addition of bioactivator has no effect on the formation of Phosphorus nutrients
 - c. The addition of bioactivators affects the formation of potassium nutrients.

Suggestion

Based on the results of the study, suggestions can be given:

1. Further research is needed on E-Colli bacteria around cattle farmers.
2. Further research is needed on the content of other micro and macro nutrients from the beginning until compost is formed.
3. Further research is needed on the biological test of the composting process of cattle solid waste.

Website : <http://jurnal.strada.ac.id/jqph> Email : jqph@strada.ac.id

REFERENCES

- Abbasi,S,A. (2013) Methane Capture From Livestock Manure. Journal Of Environmental Management 117. Hal 187
- Arikunto, Suharsini. (2010). Prosedurb Pnelitian Suatu Pendekatan Praktik. Jakarta: Rineka Cipta, hal. 211.
- Ardinal dkk (2015). Studi Potensi Pembentukan Biogas Dari Sampah Makanan Dan Ko-Substrat Fses Sapi Untuk Energi Listrik Alternatif, *Journal Litbang Industri Vol. 5 No. 2 :101-111*
- Azhar, Iman (2012). Multivariate of Variance, (Jurnal Studi Islam Madinah Vol. 7 No.1: hal 23
- Azwar, S. (2015). Dasar-Dasar Psikometrika (Edisi I I). Yogyakarta: Pustaka Pelajar.
- Basri, Elma (2017). Potensi dan Pemanfaatan Rumen Sapi Sebagai Bioaktivator, Bogor, Kementerian Pertanian RI.
- Budiman. Suyono. Fazriaty, Nadya (2018). Pengaruh Tiga Jenis Bioaktivator Ragi Terhadap Karakteristik Fisik Kompos Sampah Organik Di Rumah Sakit Cahya Kawaluyaan, *vol. 1, No.1 ISSN 2654-5411*
- Ghozali, Imam. (2016). Aplikasi Analisis Multivariate Dengan Program IBM SPSS 23 (edisi 8). Semarang : Bada Penerbit-UNDIP.

- Hidayat, Taufik. (2017, Marc 21). Sampah Membuat Jadi Kumuh dan Tidak Sehat. Propinsi Sumatera Utara. Retrieved from kotaku.pu.go.id
- Irianto, Agus. (2007). Statistik Konsep Dasar & Aplikasinya. Jakarta: Perada Media Group : hal. 271-271
- Niaga, Muhammad Syawal Akbar dkk (2020). Pengaruh Penambahan EM4 Yang Berbeda Terhadap Komposisi Hara Pupuk Organik Cair Berbahan Baku Limbah Kepala Udang Jenis *Litopenaeus vannamei*, volume 3 No. 2, ISSN : 2621-1475
- Novitasari, Devi. Caroline,Jenny. (2021). Kajian Efektivitas Pupuk Dari Berbagai Kotoran Sapi, Kambing Dan Ayam. Makalah dipresentasikan di FTSP ITATS, Surabaya.
- Nur, Thoyib dkk (2016). Pembuatan Pupuk Organik Cair dari Sampah Organik Rumah Tangga dengan Bioaktivator EM4, volume 5 No.2.e-ISSN: 2541-3481
- Nurjanah, N .dkk (2019). Pembuatan Pupuk organik Padat Dengan Aerob, *Journal of Chemical Process Engineering vol.4 No. 2 e-ISSN 2655-2967*
- Nurjasmir, Reni. Karakteristik Kompos Asal Berbagai Jenis Limbah Organik Dengan Penambahan Beberapa macam Bioaktivator. *Jurnal Ilmiah Respati Pertanian ISSN : 141-7126, Vol.2. No. Hal 619-622*
- Nurkholis. (2019). Pembuatan Pupuk Organik Padat (POP) Berbasis Bahan Kotoran Ternak Dengan Memanfaatkan Bioaktivator Isi Rumen Sapi. ISBN : 978-602-14917-8-2. Hal 170-172
- Novitasari, Devi. Caroline,Jenny. (2021). Kajian Efektivitas Pupuk Dari Berbagai Kotoran Sapi, Kambing dan Ayam. Makalah dipresentasikan di FTSP ITATS, Surabaya.
- Notoatmojo, Soekidjo. (2002). Metodologi Penelitian Kesehatan. Jakarta : Rineka Cipta, hal 46.
- Peraturan Menteri Pertanian Nomor : 02/Pert/HK.060/2/2006
- Ratrinia, Putri Wening.Uju, Pipih Suptijah (2016). Efektivitas Penambahan Bioaktivator Laut dan Limbah Cair Surimi Pada Karakteristik Pupuk Organik Cair Dari *Sargassum sp*, DOI: 10.17844/jphpi.2016.19.3.309
- SNI (2018), Pupuk Organik Padat, Jakarta : SNI 7763:2018
- Salawati, Trixie dkk (2008). Jurnal Promosi Kesehatan vol.3 No. 2
- Setiawan,Ade Iwan. (2007). Memanfaatkan Kotoran Ternak (edisi revisi). Jakarta : Penebar Swadaya, hal. 12-18
- Setiawan, Budi Susilo (2016). Bioaktivator, Beranda Dunia Tanaman
- Sharot, Siddarth. (2019). Biodiesel Production From Mutton Waste Fat –Ashort Review. *Journal Of Environmental Science and Pollution Resert* <http://doi.org/10.30799/jespr.168.19050204.page346>
- Siregar, Syofian. (2014). Statistik Parametrik Untuk Penelitian Kuantitatif. Jakarta : Bumi Aksara : hal. 77 – 90.
- Sudantha, I.M. dan Sardian(2016). Potensi Bioaktivator dan Biokompos (Mengandung Jamur *Tichrodermaspp.* dan Mikoriza) dalam Meningkatkan Kesehatan, Pertumbuhan dan Hasil Tanaman Kedelai di Lahan Kering. *Eprints.unram.ac.id*. hal 10 – 13
- Sugiyono. (2012). Metode Penelitian Pendidikan Pendekatan Kuantitatif, Kualitatif dan R & D., Bandung : Alfabeta, hal.79
- Tanzah, Ahmad. (2011). Metode Penelitian Praktis. Yogyakarta: Teras, hal. 9
- Trivana,Linda. (2017). Optimalisasi Waktu Pengomposan dan Kualitas Pupuk Kandang dari Kotoran Kambing dan Debu Sabut Kelapa Dengan Bioaktivator PROMI dan Orgadec. *JurnalSainVeterinerISSN : 2407 – 3733*. Hal 140 – 142
- Vogel. (1985). Buku Teks Analisis Anorganik Kualitatif Makro dan Semi Mikro Bagian II (edisi lima). Jakarta : PT. Kalman Media Pusaka, hal 317, 327.
- Vogel. (1985). Buku Teks Analisis Anorganik Kualitatif Makro dan Semi Mikro Bagian I (edisi lima). Jakarta : PT. Kalman Media Pusaka, hal 311
- Wellang, Rahma Musafir. Rahim, Irwan Ridwan. (2015). Study Kelayakan Kompos Menggunakan Variasi Bioaktivator. Makalah dipresentasikan di Fakultas Teknik Universitas Hasanudi, Makasar
- W.Deasy Amalia, Widiyaningrum, Priyantini (2016). Penggunaan EM4 dan MOI Limbah Tomat Sebagai Bioaktivator Pada Pembuatan Kompos, *Journal.unnes.ac.id/sju/index.php/UnnesJLifeSci, p-ISSN2252-6277*

Processing Solid Waste of Animal Beef using Bioactivators in Effort to Reduce Environmental Pollution

ORIGINALITY REPORT

10%

SIMILARITY INDEX

7%

INTERNET SOURCES

5%

PUBLICATIONS

3%

STUDENT PAPERS

PRIMARY SOURCES

- 1 Listyorini Wulandari, Indasah Indasah, Byba Melda Suhita. "Analysis of Drug Logistics Management in the Pharmacy Installation of Klaten District Health Office", Journal for Quality in Public Health, 2020
Publication 2%
- 2 Megati Hia. "EFFECT OF USING YOUTUBE ON STUDENTS' MOTIVATION IN LEARNING ENGLISH VOCABULARY", Dialectical Literature and Educational Journal, 2021
Publication 1%
- 3 fr.scribd.com
Internet Source 1%
- 4 www.nveo.org
Internet Source <1%
- 5 Gerhard Elsner, Gundula Jänsch-Kaiser, S. Gorski. "Aluminium alloys", Wiley, 2014
Publication <1%
- 6 shodhganga.inflibnet.ac.in
Internet Source <1%

7	eprints.radenfatah.ac.id Internet Source	<1 %
8	www.ijmcer.com Internet Source	<1 %
9	ijmr.net.in Internet Source	<1 %
10	garuda.kemdikbud.go.id Internet Source	<1 %
11	Azka Falaih Rizqiyana. "STEM (SCIENCE, TECHNOLOGY, ENGINEERING, AND MATHEMATICS) APPROACHES USING THEMATIC LEARNING MEDIA TO DEVELOP CRITICAL THINKING", <i>Dinamika Jurnal Ilmiah Pendidikan Dasar</i> , 2021 Publication	<1 %
12	digilibadmin.unismuh.ac.id Internet Source	<1 %
13	B. T. Shawky. "Growth and dinitrogen fixation by azospirillum, species grown under various growth conditions", <i>Acta Biotechnologica</i> , 1990 Publication	<1 %
14	Submitted to Butler University Student Paper	<1 %
15	Submitted to Universitas Brawijaya Student Paper	<1 %

16	etheses.uin-malang.ac.id Internet Source	<1 %
17	www.ajhssr.com Internet Source	<1 %
18	www.scitepress.org Internet Source	<1 %
19	ojs.uho.ac.id Internet Source	<1 %
20	repository.uinjambi.ac.id Internet Source	<1 %
21	Edy Budiman, Yonatan Parassa, Haerullah Haerullah, Dikwan Moeis, Rendra Soekarta, Muh Jamil. "The Effect of Mobile Learning Media on Student Learning Motivation in Data Structure Courses", 2018 3rd International Conference on Information Technology, Information System and Electrical Engineering (ICITISEE), 2018 Publication	<1 %
22	ejournal.unitomo.ac.id Internet Source	<1 %
23	eprints.itn.ac.id Internet Source	<1 %
24	link.springer.com Internet Source	<1 %

25

repository.unair.ac.id

Internet Source

<1 %

26

www.sciencegate.app

Internet Source

<1 %

27

digilib.unila.ac.id

Internet Source

<1 %

28

journal.lppm-unasman.ac.id

Internet Source

<1 %

29

we-didview.xyz

Internet Source

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On