

The Effect of Diabetes Self-Management Education (DSME) on Improving Self-Management and Quality of Life

by Reza Diko Utama, indasah Siti Farida Noor Layla

Submission date: 14-Apr-2022 03:50PM (UTC+0700)

Submission ID: 1810481608

File name: JQPH_VOL_4_NO_2_REZA_TERAKREDITASI_SINTA_5_NILAI_15.pdf (287.45K)

Word count: 4304

Character count: 22086

The Effect of Diabetes Self-Management Education (DSME) on Improving Self-Management and Quality of Life in Millitus Type 2 Diabetes

Reza Diko Utama¹, Indasah²,
Siti Farida Noor Layla³

¹ Health Promotion, Public Health, IIK STRADA INDONESIA

² Health Promotion, Public Health, IIK STRADA INDONESIA

³ Health Promotion, Public Health, IIK STRADA INDONESIA

Email:

rezadiko13@gmail.com

ABSTRACT

Self-management is a form of self-management or self-regulation which includes knowledge and skills in self-management. Quality of life is the level where individuals can maximize the physical and psychological functioning of life. This research method is True-Experiment with Pre-Test and Post-Test designs. The research instruments were DSMQ and WHOQOL. Independent Variables, Daiabetes Self-Management Education (DSME) Dependent Variables Self-management and quality of life. Subjects were divided into two, namely the control group totaling 55 people and the intervention group amounting to 55 people using simple random sampling technique. Data analysis using Wilcoxon and Maan-Whitney. The results of the comparison of self-management in the intervention group and the control group obtained a significance value of 0.000 less than 0.05 ($p < 0.05$) indicating that the group given the intervention was more effective at improving self-management significantly than the group that was not given the intervention. The results of the comparison of the quality of life variables of the intervention group and the control group obtained a significance value of 0.000 less than 0.05 ($p < 0.05$), which indicates that the group given the intervention was more effective in improving the quality of life significantly compared to the group that was not given the intervention.

Keywords: Education, Self-Management, Quality of Life, Diabetes Mellitus

Received : January 1st 2021

Accepted : May 3th 2021

Published : May 20th 2021

2

Copyright © 2021 IIK STRADA Indonesia
All right reserved.



This is an open-access article distributed under the terms of the Creative Commons Attribution-ShareAlike 4.0 International License.

INTRODUCTION

Non-Communicable Diseases (PNM) has become a public health problem, in recent years Non-Communicable Disease (PNM) has emerged as a key public health problem due to the high number of deaths worldwide caused by Non-Communicable Diseases (PNM), a non-communicable disease. Contagious that has received a lot of attention is Diabetes Millitus Type 2 (Pham et al., 2019). People with Type 2 Diabetes Millitus often do not have the knowledge, skills and abilities in carrying out self-management or Self-Management. Diabetes Millitus self-management behavior is very little people understand, because there are internal and external obstacles. Patients with Type 2 Diabetes Millitus still do not understand the importance of modifying self-care such as diabetes millitus diet, diabetes millitus diet, diabetes millitus physical activity, and blood sugar control are part of Diabetes Millitus self-management (Whitemore et al., 2019). Self-management is self-management of how a person can and must manage themselves so that in becoming a

successful self in their life span, someone will experience confusion about themselves, who they are and how others see themselves, someone can assess themselves multidimensional and manage themselves , 2015).

MATERIALS AND METHODS

⁸This research method is True-Experiment with Pre-Test and Post-Test designs. The research instruments were the DSMQ and WHOQOL. Independent Variables, Daiabetes Self-Management Education (DSME) Dependent Variables Self-management and quality of life. Subjects were divided into two, namely the control group totaling 55 people and the intervention group amounting to 55 people using simple random sampling technique. Data analysis using Wilcoxon and Maan-Whitney.

RESULTS

From the results of research conducted by researchers can be seen in the table as follow:

⁷**Tabel 1. Self-Management of Patients with Type II Diabetes Mellitus Comparison of Pre and Post Test in the Control Group**

Variabel		M	SD	Z	Sig.	Ket
Diabetes Diet and Diet	Pre test	1.127	0.742	-0.627	0.531	Tidak
	Post test	1.109	0.754			
Exercise / Physical Activity Diabetes	Pre test	1.025	0.767	-1.796	0.072	Tidak
	Post test	1.079	0.778			
Diabetes Blood Sugar Control	Pre test	1.167	0.766	-1.224	0.221	Tidak
	Post test	1.193	0.796			
Diabetes Health Care	Pre test	1.135	0.774	-1.721	0.085	Tidak
	Post test	1.195	0.790			
Thorough Care	Pre test	0.745	1.004	-0.322	0.748	Tidak
	Post test	0.764	0.962			
Self-Management	Pre test	1.099	0.692	-2.198	0.028	Signifikan
	Post test	1.125	0.702			

The results of the comparison of the pre-test and post-test values on the dietary pattern and diabetes diet variables in the control group obtained a significance value of 0.531 and more than 0.05 ($p > 0.05$) which indicates that there is no significant increase in diet and diabetes diet without giving it. intervention. The results of the comparison of the pre-test and post-test values on the sports / physical activity variables of diabetes in the control group obtained a significance value of 0.072 and more than 0.05 ($p > 0.05$), which indicates that there was no significant increase in sports / physical activity without giving diabetes. intervention. The results of the comparison of the pre-test and post-test values on the control variable diabetes blood sugar in the control group obtained a significance value of 0.221 and more than 0.05 ($p > 0.05$), which indicates that there was no significant increase in diabetes blood sugar control without intervention. The results of the comparison of the pre-test and post-test scores on the diabetes health care variables in the control group obtained a significance value of 0.085 and more than 0.05 ($p > 0.05$), which indicates that there was no significant increase in diabetes health care without intervention. The results of the comparison of the pre-test and post-test scores on the overall care variable in the control group obtained a significance value of 0.748 and more than 0.05 ($p > 0.05$), which indicates that there was no significant increase in overall care without intervention. The results of the comparison of the pre-test and post-test values on the self-management variable in the control group obtained a significance value of 0.028 and less than 0.05 ($p < 0.05$) which indicates a small amount of improvement in self-management.

5

Tabel 2. Quality of Life for Patients with Type II Diabetes Mellitus Comparison of Pre-Test and Post in the Control Group

Variabel		M	SD	Z	Asymp Sig.	Significancy
Physical Health	Pre test	1.979	0.996	-1.601	0.109	No
	Post test	2.012	0.981			
Psychology	Pre test	2.185	0.985	-0.759	0.448	No
	Post test	2.204	1.012			
Social	Pre test	2.280	0.924	-1.341	0.180	No
	Post test	2.240	0.922			
Environment	Pre test	1.979	0.892	-1.493	0.135	No
	Post test	1.943	0.928			
Thorough	Pre test	2.336	1.093	-0.791	0.429	No
	Post test	2.364	1.120			
Quality of Life	Pre test	2.099	0.892	-0.169	0.866	No
	Post test	2.099	0.908			

The results of the comparison of the pre-test and post-test scores on the physical health variables in the control group obtained a significance value of 0.109 and more than 0.05 ($p > 0.05$), which indicates that there was no significant increase in physical health without intervention. The results of the comparison of the pre-test and post-test scores on psychological variables in the control group obtained a significance value of 0.448 and more than 0.05 ($p > 0.05$) which indicates that there is no significant psychological improvement without the intervention. The results of the comparison of the pre-test and post-test scores on social variables in the control group obtained a significance value of 0.180 and more than 0.05 ($p > 0.05$), which indicates no significant social improvement without intervention. The results of the comparison of the pre-test and post-test values for environmental variables in the control group obtained a significance value of 0.135 and more than 0.05 ($p > 0.05$), which indicates that there is no significant environmental improvement without intervention. The results of the comparison of the pre-test and post-test scores on the overall variables in the control group obtained a significance value of 0.429 and more than 0.05 ($p > 0.05$), which indicates that there was no significant overall increase without intervention. The results of the comparison of the pre-test and post-test scores on the quality of life variable in the control group obtained a significance value of 0.866 and more than 0.05 ($p > 0.05$), which indicates that there was no significant increase in quality of life without intervention.

7

Tabel 3. Self-Management of Patients with Type II Diabetes Mellitus Comparison of Pre and Post Test in the Intervention Group

Variabel		M	SD	Z	Asymp Sig.	Significancy
Diabetes Diet and Diet	Pre test	1.477	0.775	-5.561	0.000	Sig
	Post test	2.491	0.821			
Exercise / Physical Activity Diabetes	Pre test	1.431	0.921	-5.226	0.000	Sig
	Post test	2.485	0.812			
Diabetes Blood Sugar Control	Pre test	1.367	0.773	-5.860	0.000	Sig
	Post test	2.458	0.900			
Diabetes Health Care	Pre test	1.418	0.792	-5.845	0.000	Sig
	Post test	2.539	0.803			
Thorough Care	Pre test	1.436	0.996	-4.983	0.000	Sig
	Post test	2.491	0.979			
Self-Management	Pre test	1.421	0.734	-6.128	0.000	Sig
	Post test	2.490	0.820			

The results of the comparison of the pre-test and post-test values on the variable dietary diet and diabetes diet in the intervention group obtained a significance value of 0.000 and less than 0.05 ($p < 0.05$) which indicates a significant increase in dietary patterns and diabetic diet with the intervention. The results of the comparison of the pre-test and post-test values on the sports / physical activity variables of diabetes in the intervention group obtained a significance value of

0.000 and less than 0.05 ($p < 0.05$) which indicates a significant increase in sports / physical activity with diabetes with the intervention. The results of the comparison of the pre-test and post-test values on the control variable diabetes blood sugar in the intervention group obtained a significance value of 0.000 and less than 0.05 ($p < 0.05$) which indicates a significant increase in diabetes blood sugar control with the intervention. The results of the comparison of the pre-test and post-test values on the diabetes health care variable in the intervention group obtained a significance value of 0.000 and less than 0.05 ($p < 0.05$) which indicates a significant increase in diabetes health care with the provision of intervention. The results of the comparison of the pre-test and post-test scores on the overall care variable in the intervention group obtained a significance value of 0.000 and less than 0.05 ($p < 0.05$) which indicates a significant increase in overall care with the intervention. The results of the comparison of the pre-test and post-test scores on the self-management variable in the intervention group obtained a significance value of 0.000 and less than 0.05 ($p < 0.05$) which indicates a significant increase in self-management with the provision of intervention

Tabel 4. Quality of Life for Patients with Type II Diabetes Mellitus Comparison of Pre and Post Test in the Intervention Group

Variabel		M	SD	Z	Asymp Sig.	Significancy
Physical Health	Pre test	2.148	1.280	-5.689	0.000	Sig
	Post test	3.720	1.477			
Psychology	Pre test	2.429	0.952	-5.831	0.000	Sig
	Post test	3.655	1.182			
Social	Pre test	2.385	1.160	-5.552	0.000	Sig
	Post test	3.273	1.021			
Environment	Pre test	2.205	1.063	-4.893	0.000	Sig
	Post test	2.780	1.072			
Thorough	Pre test	2.727	1.018	-5.404	0.000	Sig
	Post test	3.964	1.258			
Quality of Life	Pre test	2.331	1.062	-5.952	0.000	Sig
	Post test	3.392	1.165			

The results of the comparison of the pre-test and post-test values on the physical health variables in the intervention group obtained a significance value of 0.000 and less than 0.05 ($p < 0.05$), which indicates a significant increase in physical health with the intervention. The results of the comparison of the pre-test and post-test scores on psychological variables in the intervention group obtained a significance value of 0.000 and less than 0.05 ($p < 0.05$) which indicates a significant psychological improvement with the intervention. The results of the comparison of the pre-test and post-test scores on social variables in the intervention group obtained a significance value of 0.000 and less than 0.05 ($p < 0.05$), which indicates a significant social improvement with the intervention. The results of the comparison of the pre-test and post-test scores on environmental variables in the intervention group obtained a significance value of 0.000 and less than 0.05 ($p < 0.05$) which indicates a significant increase in the environment with the provision of intervention. The results of the comparison of the pre-test and post-test scores on the overall variables in the intervention group obtained a significance value of 0.000 and less than 0.05 ($p < 0.05$) which indicates a significant overall increase with the provision of the intervention. The results of the comparison of the pre-test and post-test scores on the quality of life variable in the intervention group obtained a significance value of 0.000 and less than 0.05 ($p < 0.05$) which indicates a significant increase in the quality of life with the intervention.

7
Table 5. Self-Management of Patients with Type II Diabetes Mellitus Comparison of Control and Intervention Groups

Variabel		M	SD	Z	Asymp Sig.	Significancy
Diabetes Diet and Diet	Intervensi	2.491	0.821	-6.763	0.000	Sig
	Kontrol	1.109	0.754			
Exercise / Physical Activity Diabetes	Intervensi	2.485	0.812	-6.732	0.000	Sig
	Kontrol	1.079	0.778			
Diabetes Blood Sugar Control	Intervensi	2.458	0.900	-6.231	0.000	Sig
	Kontrol	1.193	0.796			
Diabetes Health Care	Intervensi	2.539	0.803	-6.986	0.000	Sig
	Kontrol	1.195	0.790			
Thorough Care	Intervensi	2.491	0.979	-6.824	0.000	Sig
	Kontrol	0.764	0.962			
Self-Management	Intervensi	2.490	0.820	-7.430	0.000	Sig
	Kontrol	1.125	0.702			

The results of the comparison of the variable dietary diet and diabetes diet between the intervention group and the control group obtained a significance value of 0.000 and less than 0.05 ($p < 0.05$), which indicates that the group given the intervention was more effective in improving the diabetes diet and diet significantly. compared with the group not given the intervention. The results of the comparison of the diabetes exercise / physical activity variable between the intervention group and the control group obtained a significance value of 0.000 and less than 0.05 ($p < 0.05$), which indicates that the group given the intervention was more effective at increasing diabetes exercise / physical activity significantly. compared with the group not given the intervention. The results of the comparison of the diabetes blood sugar control variable between the intervention group and the control group obtained a significance value of 0.000 and less than 0.05 ($p < 0.05$), which indicates that the group given the intervention was more effective at improving diabetes blood sugar control significantly compared to the group that was not given the intervention. on the diabetes health care variable between the intervention group and the control group, a significance value of 0.000 and less than 0.05 ($p < 0.05$) was obtained, which indicates that the group given the intervention was more effective at improving diabetes health care significantly compared to the group that was not given intervention. The results of the comparison of the post-test scores on the overall care variable between the intervention group and the control group obtained a significance value of 0.000 and less than 0.05 ($p < 0.05$), which indicates that the group given the intervention was more effective at increasing overall care significantly compared to the who were not given the intervention. The results of the comparison of the self-management variable between the intervention group and the control group obtained a significance value of 0.000 and less than 0.05 ($p < 0.05$), which indicates that the group given the intervention was more effective at improving self-management significantly compared to the group that was not given. intervention.

8
Table 6. Quality of Life of Patients with Type II Diabetes Mellitus Comparison in Control and Intervention Group

Variabel		M	SD	Z	Asymp Sig.	Significancy
Physical Health	Intervensi	3.720	1.477	-5.272	0.000	Sig
	Kontrol	2.012	0.981			
Psychology	Intervensi	3.655	1.182	-5.661	0.000	Sig
	Kontrol	2.204	1.012			
Social	Intervensi	3.273	1.021	-5.239	0.000	Sig
	Kontrol	2.240	0.922			
Environment	Intervensi	2.780	1.072	-4.528	0.000	Sig
	Kontrol	1.943	0.928			
Thorough	Intervensi	3.964	1.258	-5.925	0.000	Sig
	Kontrol	2.364	1.120			
Quality of Life	Intervensi	3.392	1.165	-5.236	0.000	Sig
	Kontrol	2.099	0.908			

The results of the comparison of the physical health variables between the intervention group and the control group obtained a significance value of 0.000 and less than 0.05 ($p < 0.05$), which indicates that the group given the intervention was more effective at improving physical health significantly than the group that was not given intervention. The results of the comparison of the psychological variables between the intervention group and the control group obtained a significance value of 0.000 and less than 0.05 ($p < 0.05$), which indicates that the group given the intervention was more effective at improving psychology significantly than the group that was not given the intervention. The results of the comparison of the social variables between the intervention group and the control group obtained a significance value of 0.000 and less than 0.05 ($p < 0.05$), which indicates that the group given the intervention was more effective at improving social significantly than the group that was not given the intervention. The results of the comparison of the environmental variables between the intervention group and the control group obtained a significance value of 0.000 and less than 0.05 ($p < 0.05$), which indicates that the group given the intervention was more effective in improving the environment significantly than the group that was not given the intervention. The results of the comparison of the overall variable between the intervention group and the control group obtained a significance value of 0.000 and less than 0.05 ($p < 0.05$), which indicates that the group that was given the intervention was more effective at increasing overall significantly compared to the group that was not given the intervention. The results of the comparison of the quality of life variable between the intervention group and the control group obtained a significance value of 0.000 and less than 0.05 ($p < 0.05$), which indicates that the group given the intervention was more effective at improving quality of life significantly compared to the group that was not given intervention.

DISCUSSION

According to Severtina, H., (2019), said that health education is very influential on the level of knowledge, besides that health education can affect the level of knowledge so that it can have changes in healthy behavior. According to Bekele, B. B., (2020), said that Diabetes Self-Management Education (DSME) has a positive impact on the level of self-management and quality of life in people with Diabetes Mellitus and can reduce the risk of complications in Diabetes Mellitus. Diabetes Self-Management Education (DSME) is a process of facilitating skills, knowledge and abilities in self-care or diabetes mellitus self-management. According to Agustiningrum, R. (2019), in his research, he said that Diabetes Self-Management Education (DSME) is very effective for self-care in Diabetes Mellitus patients. From the results of research conducted by researchers in the intervention group, Diabetes Self-Management Education (DSME) is a Diabetes Mellitus health education method that is effective in increasing knowledge. Skills and abilities in self-management and quality of life in people with Type II Diabetes Mellitus. From the results of research conducted by researchers in the intervention group, it is proven that they have knowledge, skills and abilities in self-management and experience improvement in self-management and quality of life after being given the Diabetes Self-Management Education (DSME) health education method or program. Diabetes Self-Management Education (DSME) was able to improve self-management and quality of life in the intervention group so that there were significant changes in the level of self-management and quality of life in people with Type II Diabetes Mellitus. According to Maulana (2019), one of the strategies in obtaining behavior change in people with Diabetes Mellitus is by providing health information or education. The process of empowering or empowering people with Diabetes Mellitus can be done by providing information or education such as health education in changing behavior and attitudes in order to achieve self-management and a good quality of life. According to Notoadmodjo (2007), the level of education greatly determines a person's ability to understand the knowledge obtained, such as the higher the level of one's education, the more someone is willing to accept and understand information. Research conducted by Wiastuti (2017) states that high levels of education will tend to easily absorb health information so that it can change a person's behavior. According to Haas, (2014) in his journal entitled National Standards DSME is a process to facilitate the knowledge, skills, and abilities needed for diabetes self-care (Self-Management). This process combines the needs, goals, and life experiences of people with diabetes or prediabetes and is guided by evidence-based standards. The overall goal of DSME is to support decision making, self-care behavior, problem solving, and active collaboration with health care teams and to improve clinical outcomes, health status, and quality of life.

CONCLUSION

Based on the results of the research that has been presented, Diabetes Self-Management Education (DSME) is proven to be effective in improving self-management and quality of life in Diabetes Mellitus sufferers, Diabetes Mellitus sufferers are able to determine the right nutritional diet, be able to exercise / physical activity, control blood sugar levels and being able to perform Diabetes Mellitus health so that the treatment of Diabetes Mellitus sufferers who have good levels of self-management and good quality of life. Diabetes Self-Management Education (DSME) can improve self-management and quality of life in people with Type II Diabetes Mellitus. The results of the study showed that the intervention group that was given treatment or intervention proved to be very effective in improving self-management and quality of life and was inversely proportional to the control group that was not given treatment.

ACKNOWLEDGMENTS


Researchers are grateful to all respondents who have been cooperative in participating in all research activities. Researchers and thanks to the related parties of the Kempo Health Center, Kempo District, Dompu Regency who have allowed researchers to carry out the process of research activities

CONFLICTS OF INTEREST

In the research that has been done, this research does not have a conflict of interest from other parties.

REFERENCES

- Agustiningrum, R., Kusbaryanto, K., Keperawatan, M., & Kedokteran, F. (2019). *Efektifitas Diabetes Self Management Education Terhadap Self Care Penderita Diabetes Mellitus: A Literature Review*. 6(2), 558–563
- Bekele, B. B., Negash, S., Bogale, B., Tesfaye, M., & Getachew, D. (2020). *The effectiveness of diabetes self-management education (DSME) on glycemic control among T2DM patients randomized control trial : systematic review and meta-analysis protocol*
- Maulana, H.DJ. (2019). *Promosi Kesehatan*. Jakarta: EGC
- Haas. (2014). *National Standar For Diabetes Self-Management Education of Support.Diabetes Care*
- Notoatmodjo, (2010). *Prilaku Kesehatan*. Jakarta: PT. Rineka Cipt
- Pham, B. D., Kim, B. G., Nguyen, T. T. H., & Hoang, V. M. (2019). Exposure to messages on risk factors for noncommunicable diseases in a rural Province of Vietnam. *BioMed Research International*, 2019. <https://doi.org/10.1155/2019/7962947>
- Seventina, H., Tinggi, S., Kesehatan, I., Astrid, M., Concentration, M. S., Indonesia, J., Susilo, W. H., & Lecture, R. (2020). *Effect of Health Education and Diabetic Foot Exercise on Sensory Perception and Knowledge level of Patients With Type 2 Diabetes Mellitus in Cirebon District General Hospital in West Java Province*. 27(ICoSHEET 2019), 362–365.
- Whittemore, R., Vilar-Compte, M., De La Cerda, S., Marron, D., Conover, R., Delvy, R., Lozano-Marrufo, A., & Pérez-Escamilla, R. (2019). Challenges to diabetes self-management for adults with type 2 diabetes in low-resource settings in Mexico City: A qualitative descriptive study. *International Journal for Equity in Health*, 18(1), 1–10. <https://doi.org/10.1186/s12939-019-1035-x>
- Wiastruti Mariana Siti Rondhianto, Nur Widayanti. (2017). *Pengaruh Diabetes Self-Management Education and Support (DSME/S) Terhadap Stres Pasien Diabetes Mellitus Tipe II di Wilayah Kerja Puskesmas Patrang Kbaupaten Jember*. Universitas Jember.Jember



The Effect of Diabetes Self-Management Education (DSME) on Improving Self-Management and Quality of Life

ORIGINALITY REPORT

7%

SIMILARITY INDEX

4%

INTERNET SOURCES

4%

PUBLICATIONS

3%

STUDENT PAPERS

PRIMARY SOURCES

- 1** Submitted to University of Missouri, Kansas City
Student Paper 1%
- 2** Moses Lamere, Ratna Wardani. "Analysis of Nurse Characteristics with Work Motivation in Wamena Hospital", Journal for Quality in Public Health, 2021
Publication 1%
- 3** Submitted to Prince of Songkla University
Student Paper 1%
- 4** Undatun Ni'mah, Nia Sari, Yuly Peristiowati. "Analyse the Role of Cadre, Parenting and Food Intake to Nutrition Status of Toddler", Journal for Quality in Public Health, 2019
Publication 1%
- 5** munin.uit.no
Internet Source 1%
- 6** Submitted to University of Sheffield
Student Paper 1%

7	Lenni Sastra, Ise Reni. "The Effect of Indonesian Traditional Music on Stress in Type II Diabetes Mellitus Patients", KnE Life Sciences, 2022 Publication	1 %
8	journal.unnes.ac.id Internet Source	<1 %
9	Oluwadamilare Akingbade, Khanh Thi Nguyen, Ka Ming Chow. "Effect of mHealth interventions on psychological issues experienced by women diagnosed with breast cancer receiving chemotherapy: A systematic review and meta-analysis", Cold Spring Harbor Laboratory, 2022 Publication	<1 %
10	clinmedjournals.org Internet Source	<1 %
11	ir.lib.ruh.ac.lk Internet Source	<1 %
12	Submitted to Institute of Health & Management Pty Ltd Student Paper	<1 %
13	www.coursehero.com Internet Source	<1 %
14	www.jqph.org Internet Source	<1 %

15 www.hindawi.com Internet Source <1 %

16 scholarworks.waldenu.edu Internet Source <1 %

17 Erni Tri Indarti, Lexy Oktora Wilda, Yiyin Nuvitasari. "Water Boiling Celery Affects Blood Pressure in Elderly with Hypertension", *Journal for Quality in Public Health*, 2020
Publication <1 %

18 Risa Nurhayati, Indasah Indasah, Byba Melda Suhita. "Family Support in Effort Reduce Hospitalization Reaction in Children of Preschool in Angrek Room Nganjuk Hospital", *JOURNAL FOR QUALITY IN PUBLIC HEALTH*, 2018
Publication <1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On