

The Effect Consumption Of Red Beans (*Phaseolus Vulgaris* L) Boiled On Hemoglobin Levels

by Candra W, Shanty N, Miftakhur R, Retno P Y, Nita Anggrawati Wulandari, Riza Tsalatsatul M, Yeni P

Submission date: 31-May-2021 05:32PM (UTC+0700)

Submission ID: 1597747456

File name: EJMCM_Volume_7_Issue_2_Pages_4754-4759.pdf (312.8K)

Word count: 2539

Character count: 13821

The Effect Consumption Of Red Beans (*Phaseolus Vulgaris L*) Boiled On Hemoglobin Levels In Adolescent In Institute Of Health Science STRADA Indonesia

Candra Wahyuni, Shanty Natalia, Miftakhur Rohmah, Retno Palupi Yoni Siwi, Nita Dwi Astikasari, Anggrawati Wulandari, Riza Tsalatsatul Mufida, Yeni Puspitasari

Department of Midwifery, Institute of Health Science Strada Indonesia Kediri, East Java, Indonesia

Corresponding author: candrawahyuni85@gmail.com

Abstract: Anemia in general is a condition where hemoglobin and erythrocyte levels are lower than normal. Anemia that is occurred by many adolescents due to Fe deficiency and can be called Fe deficiency anemia (Tarwoto, 2014). The purpose of this study was to determine the effect of consumption of red beans (*Phaseolus Vulgaris L*) boiled on hemoglobin levels in adolescent in institute of health science STRADA Kediri. The research design used was pre-experiment with one group pre-post test design. The population in this study were all third semester students in Institute of health Science STRADA who experienced anemia with a sample of 18 respondents. The sampling technique used was Purposive sampling. The results showed that all (100%) had mild anemia before treatment, after the treatment, the results of all respondents (100%) experienced a change to non-anemia. Based on the results statistically using test, Wilcoxon the obtained p value $0.005 < \alpha 0.05$ then H_0 is rejected and H_1 accepted, which means there is the effect of red beans (*PhaseolusvulgarisL*) boil on hemoglobin levels in adolescent in institute of health science STRADA Kediri. It can be concluded that boiled red beans (*Phaseolus Vulgaris L*) contain iron which is useful for increasing hemoglobin levels in the blood. It is recommended for respondents to use boiled red beans as food to prevent anemia.

Keywords--- Red beans, Hemoglobin levels, Adolescent

I. INTRODUCTION

Adolescent anemia is more common than children and adults, especially for adolescent girls. This is because teenage girls experience menstruation, low food intake, increased iron requirements due to the process of *growth spurt* and limiting to eat so that the body deficiencies nutritional such as iron [2]. Anemia in general condition where hemoglobin and erythrocyte levels are lower than normal. Iron deficiency anemia itself is a state of decreased iron concentration in the body, both in storage, circulation and in the form of bonds with heme so that it can cause a decrease in red blood cell concentration. Iron deficiency anemia is a stage of severe iron deficiency [22][31].

Based on WHO data (2014) organization states that anemia affects 1.62 billion people in the world (24.8%). Based on Basic Health Research [18]. It is known that the prevalence of anemia in adolescents aged 15-24 years is 18.4% and in the fertile age group of women is 16.9%, so it can also be categorized as moderate health problems. Data from the East Java Provincial Health Office in 2017 states that 50-60% of young women in East Java suffer from anemia or red blood deficiency. From Riskesdas, the prevalence of national nutritional anemia among adolescents 13-18 years was 22.7% [25]. Based on the results of a preliminary study conducted on March 29, 2019 at Institute of health science STRADA, a midwifery midwifery study program student, it was found that 7 (70%) of 10 female students were anemic, 4 (40%) were due to sleep patterns and 3 (30%) people because of unhealthy eating patterns.

The prevalence of anemia in 2017 is caused by various factors such as economic, socio-cultural and the direct cause is an imbalance between food intake and nutritional needs. Anemia can be caused by a lack of food sources that contain iron, because iron is an important compound as a constituent of hemoglobin and this occurs because of poor dietary care, irregular and does not balance the nutritional adequacy needed [22][28]. The symptoms that often arise include dizziness, weakness, fatigue, fatigue and lethargy. Sometimes anemia does not cause obvious symptoms such as fatigue when exercising, difficulty concentrating and easily forgetting. The unfortunate situation is that most sufferers do not know and do not realize it. In general, someone has begun to suspect anemia if the condition is getting worse, so that the symptoms are more obvious, such as pale skin, palpitations, dizziness, easy to run out of breath when climbing stairs and exercise. With the occurrence of anemia in adolescents can have an impact on decreased work productivity or academic ability in school, due to the lack of enthusiasm for learning and concentration. Anemia can disturb growth where height and weight become imperfect. In addition, the immune system will decrease so it is susceptible to disease [13][29].

Anemia can be overcome by means of pharmacological therapy and non-pharmacological therapy. Pharmacological therapy to overcome anemia is by consuming blood booster tablets or Fe tablets. While non-pharmacological therapy to overcome anemia is to consume foods that are high in iron content. Foods that contain lots of iron are chicken, dried fruits, egg yolks, lean meat, spinach, liver, and beans. Nuts that contain lots of iron include soybeans, green beans and red beans. Red beans have a content that can overcome anemia, the content is iron (Fe). The iron content in red beans can help control the amount of hemoglobin in the blood [9][30].

II. METODE

This study included a *Pre Experiment* with approach *one group pre posttest control group with design* subjects were assessed before being treated and after being treated. Based on the scope of the study it was type of *inferential* research, based on the place of research it was type of field research, based on the presence or absence of treatment it was *experimental*, based on data collection it was observational research and based on the data source it was primary research design. Statistical test results using *Wilcoxon test*.

The population in this study were 37 semester students of D III Midwifery Program. Technique sampling used was "*purposive sampling*". The sample in this study were 18 students on the semester III of Midwifery program who experience anemia. The instrument used in this study was the *observation sheet*, all respondents observed hemoglobin levels before and after treatment. Checking Hemoglobin levels of anemia with HB elektrik / Haemometer.

The research procedures by measuring level hemoglobin before giving red beans, then given boiled kidney beans that is 450 grams per day for 3 days and measuring level hemoglobin after giving red beans [4]. Before study, researchers follow procedures ethical clearance on campus and have obtained a certificate of ethics, then conducted research. Before the study, the researcher explained the research procedure to prospective respondents and filled out the inform.consent.

III. RESULTS

- Univariate Analysis

Table 1. Frequency Distribution Characteristics of Respondents by Age in Institute of health science STRADA

Age	n	%
Early Adolescents	0	0
Mid adolescents	0	0
Late Adolescents	18	100
Total	18	100

Based on table 1. above can be interpreted that all of them are late teens (19-21) years, that is 18 respondents (100%).

- Analysis of Bivariate

Table 2. Frequency Distribution Based on Hemoglobin Levels Before giving Red Beans (*Phaseolus Vulgaris L*) Boil in Adolescent Girls in Institute of health science STRADA

Before being given red beans (<i>Phaseolus Vulgaris L</i>)	n	%
Severe Anemia	0	0
Moderate Anemia	0	0
Mild Anemia	18	100
No Anemia	0	0
Total	18	100

Based on table 2 above, it can be interpreted that all of them have mild anemia which is 18 respondents (100%).

Table 3. Frequency Distribution Based on Hemoglobin Levels after Being Given Red Beans (*Phaseolus Vulgaris L*) Boil in Adolescent Girls in Institute of health science STRADA

After Given Red Beans (<i>Phaseolus Vulgaris L</i>)	n	%
Severe Anemia	0	0
Moderate Anemia	0	0
Anemia Mild	0	0
No Anemia	18	100
Total	18	100

Based on table 3 above, it can be interpreted that all of them did not experience anemia as many as 18 respondents (100%).

IV. DISCUSSION

Based on table 3 above it can be interpreted that all (100%) experienced a change from mild anemia to not anemia, as many as 18 respondents. Based on the statistical test results

using *Wilcoxon test*, it can be interpreted that the $p\text{-value} < \alpha$ is $0.005 < 0.05$ so that H_0 is rejected and H_1 is accepted, meaning that there is an effect of giving red beans (*Phaseolus Vulgaris L*) Boiled on hemoglobin levels in adolescent in institute of health science Strada Indonesia.

The result above is suitable with the results of research conducted by Faridah and Indraswari [7] which states that consuming nuts can help increase hemoglobin levels in adolescent girls. Anemia that occurs in adolescents can occur due to consumption patterns of Indonesian people who are still dominated by vegetables as a source of iron (non-heme iron). In addition, the cause of iron deficiency anemia is influenced by increased body needs, due to suffering from chronic disease, blood loss due to menstruation and parasitic infections [17].

Boiled red beans have a variety of ingredients that are beneficial to health, one of which is the content of iron or Fe [4]. The content of iron in boiled red beans can help increase hemoglobin levels in the body. The mechanism of iron can help increase hemoglobin levels in the body by means of biochemical reactions in the body, from the biochemical process iron is able to produce hemoglobin [8]. Biochemical reactions in the form of inorganic complexes Fe^{3+} are broken down during digestion and some are converted from Fe^{3+} to Fe^{2+} which is more readily absorbed. The conversion of Fe^{3+} to Fe^{2+} is facilitated by the presence of endogenous factors such as HCl in gastric secretory fluids, components of nutrients derived from food such as vitamin C or meat and fish. This mechanism plays a role in increasing hemoglobin in the body [17].

Lack of protein intake will result in obstructed transportation so that iron deficiency will occur. Absorption of substances that occur in the small intestine is carried out by means of protein transport namely transferrin and ferritin. Iron-containing transfer which functions to transport iron into the bone marrow to form hemoglobin. Many protein sources are found in spinach, liver, fruit, nuts. Protein is important for DNA synthesis. one deficiency can increase cell regeneration and cause anemia, the size of red blood cells is larger than normal [26].

Based on the results of the research, researchers believe that the majority of respondents experienced a change from mild anemia to not anemia. Changes in hemoglobin levels occur because of the iron content in boiled red beans. The content of iron in boiled red beans can help increase hemoglobin levels in the body. The mechanism of iron can help increase hemoglobin levels in the body by means of biochemical reactions in the body, from the biochemical process iron is able to produce hemoglobin.

V. CONCLUSION

Conclusion this research is boiled red beans (*Phaseolus Vulgaris L*) can increase hemoglobin levels in young women because one of which is the content of iron or Fe. Hemoglobin levels are also influenced by many factors, nutrition status, activity, stressors, illness etc.

VI. CONFLICT OF INTEREST

No conflicts of interest have been declared.

VII. ACKNOWLEDGEMENTS

Authors of this study would like to thank the Faculty of Nursing for providing the opportunity to present this study.

REFERENCES

- [1] Almaitzer. (2014). *Prinsip Dasar Ilmu Gizi*. Jakarta: Gramedia Pustaka Utama.
- [2] Arisman. (2011). *Gizi Dalam Daur Kehidupan*. Jakarta: EGC.
- [3] Asrinah. (2015). *Asuhan Kebidanan Masa Persalinan*. Yogyakarta: Graha Ilmu.
- [4] Astawan, M. (2014). *Sehat dengan Hidangan Kacang dan Biji-bijian*. Jakarta: Penebar Swadaya.
- [5] Badriah. (2016). *Gizi dalam Kesehatan Reproduksi*. Bandung: Refika Aditama.
- [6] Benoist, B., Mclean, E., Egli, I., & Cogswell, M. (2013). *World Wide Prevalance of Anemia*. WHO Global Database on Anemia Geneva: World Health Organization.
- [7] Faridah, U., & Indraswari, V. (2017). Pemberian kacang hijau sebagai upaya peningkatan kadar hemoglobin pada remaja putri. *THE 5TH URECOL PROCEEDING*.
- [8] Handayani, & Hariwibowo. (2012). *Asuhan Keperawatan Pada Klien Dengan Gangguan Sistem Hematologi*. Jakarta: Salemba Medika.
- [9] Hanifah, T. (2012). *Perawatan Antenatal dan Peranan Asam Folat dalam Upaya Meningkatkan Kesejahteraan Ibu Hamil dan Janin*. Medan: USU.
- [10] Haniva-Kun. (2016). Kadar Hemoglobin Menurut WHO. *Tips Solusi Kesehatan*.
- [11] Indonesia, D. K. (2010). *Program Penanggulangan Anemia Gizi Pada Wanita Usia Subur*. Depkes RI.
- [12] Istiany, Ari, & Rusilanti. (2013). *Gizi Terapan*. Jakarta: Remaja Rosdakarya.
- [13] Kediri, P. K. (2015). *Profil Kesehatan Kota Kediri*. Kediri.
- [14] Manuaba. (2015). *Ilmu Kebidanan Penyakit Kandungan dan KB*. Jakarta: EGC.
- [15] Novi, A. (2016) Kacang Merah Rebus.
- [16] Permaesih, D., & Herman, S. (2015). Faktor-faktor yang mempengaruhi Anemia Pada Remaja. *Penelitian Kesehatan*.
- [17] Proverawati, K. (2013). *Buku Ajar Gizi untuk Kebidanan*. Yogyakarta: Nuha Medika.
- [18] Riskesdas, R. K. (2016). *Badan Penelitian dan Pengembangan Kesehatan Kementerian RI tahun 2016*. Jakarta:Depkes.
- [19] Rukmana, R. (2012). *Usaha Tani*. Yogyakarta: Kanisius.
- [20] Sugiyono. (2014). *Metode Penelitian Kuantitatif, Kualitatif dan R&D*. Bandung: Alfabeta.

- [21] Swara, T. (2014). *Suplementasi Iron Zinc Antisipasi Anemia Remaja Putri*. Pekanbaru.
- [22] Tarwoto. (2014). *Buku Saku Anemia Pada Ibu Hamil, Konsep dan Penatalaksanakannya*. Jakarta: Trans info Medika.
- [23] Varney, H. (2007). *Buku Ajar Asuhan Kebidanan*. Jakarta: EGC.
- [24] Waryana. (2015). *Gizi Reproduksi*. Yogyakarta: Pustaka Rihama.
- [25] Jenny Anna Siauta. (2020). *Journal For Quality in Women's Health. Vol 3 No. 1 Maret 2020*.
- [26] JOMIS (Journal Of Midwifery Science) P-ISSN : 2549 – 2543 E-ISSN : 2579 - 7077 Vol 4. No. 1 Januari 2020
- [27] A. Worku dan O. Sahu. "Significance Of Fermentation Process On Biochemical Properties Of Phaseolus Vulgaris (Red Beans). " *Biothechnol Rep.*, Vol. 16, hlm. 5 – 11, Des 2017.
- [28] Aldahmani, S., Al-Shami, S. A., Adil, H., & Sidek, S. (2020). A review paper on moocs development stages, types, and opportunities and challenges. *Systematic Reviews in Pharmacy*, 11(12), 172–179. <https://doi.org/10.31838/srp.2020.12.28>
- [29] Ali, M., Halim, A., Yunita, R., Priyanka, W., Permana, N., Arief, B. B., Diponegoro, U., & Indonesia, U. I. (2020). A PSYCHOLOGICAL PERSPECTIVE OF CORPORATE. 11(12), 925–930.
- [30] Ali, R. (2020). Survey Interactive Synthesis of Animation Model Based on Tracking User ' s Utterance and Facial Expression. 11(12), 907–915.
- [31] ALkamil, R. D. S., Al-Bahadily, D. C. H., Chalooob, R., Shari, F. H., & AL-Salman, H. N. K. (2020). Estimation of truffles extracts and study the biological efficacy of ethyl 6-methyl-2-oxo-4-(2-thienyl)-1,2,3,4-tetrahydropyrimidine-5-carboxylate (EMOTTC) as one of the extracts against ophthalmic bacteria. *Systematic Reviews in Pharmacy*, 11(12), 78–87. <https://doi.org/10.31838/srp.2020.12.13>

A. Worku dan O. Sahu, "Significance of fermentation process on biochemical properties of Phaseolus vulgaris (red beans)," *Biotechnol. Rep.*, vol. 16, hlm. 5–11, Des 2017

APPENDIX

Table 1. Standard Operational Procedures Consumption Of Red Beans Boiled [27]

Item of SOP		
Ingredient	Red Beans	150 gr
	Water	500 ml
Countains	Every 150 gr red beans :	
	water	14.40 %
	ash	2.19 %
	protein	2.23 %
	fat	2.10 %

	carbohydrate	79.09 %
Process	Boil red beans in boiling water until softer or about 15 - 20 minutes	
Dosage	Take 150 gr red beans boiled, repeat tree times in a-24 hours. Giving red beans boiled for three days	

The Effect Consumption Of Red Beans (Phaseolus Vulgaris L) Boiled On Hemoglobin Levels

ORIGINALITY REPORT

12%

SIMILARITY INDEX

6%

INTERNET SOURCES

4%

PUBLICATIONS

9%

STUDENT PAPERS

PRIMARY SOURCES

1	Submitted to Universiti Malaysia Sabah Student Paper	4%
2	Submitted to Rogers State University Student Paper	2%
3	Submitted to Chulalongkorn University Student Paper	1%
4	Submitted to La Trobe University Student Paper	1%
5	journal.unj.ac.id Internet Source	1%
6	Andika Bagus Nur Rahma Putra, Amat Mukhadis, Sumarli Sumarli, Eddy Sutadji, Poppy Puspitasari, Mahfudi Sahly Subandi. "Innovation of smart-curriculum model through campus-school-industry synchronization for vocational learning in the era of education 4.0", Proceedings of the 5th International Conference on Sustainable	1%

Information Engineering and Technology, 2020

Publication

7	text-id.123dok.com Internet Source	1 %
8	Submitted to Universitas Pendidikan Ganesha Student Paper	1 %
9	ejournalnwu.unw.ac.id Internet Source	<1 %
10	Trias Mahmudiono, Stefania Widya Setyaningtyas, Ilham Akhsanu Ridlo, Qonita Rachmah, Triska Susila Nindya. "Consumption of Iron Supplement and Anemia Among Indonesian Adolescent Girls", Global Journal of Health Science, 2019 Publication	<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On